Author Index

Alexander, P. W., Głód, B. K. and Haddad, P. R.
Mixed potential response at metallic electrodes in potentiometric detection for ion chromatography 589(1992)201

Alexander, P. W., see Głód, B. K. 589(1992)209

Alhama, J., López-Barea, J., Toribio, F. and Roldán, J. M.
Purification and determination of glutamine synthetase by high-performance immunoaffinity chromatography 589(1992)121

An, D., see Zhou, W. 589(1992)358

Anelli, P. L., Tomba, C. and Uggeri, F.
Optical resolution of 2-chloro-3-phenylmethoxypropanoic acid after derivatization with (*S*)-2-octanol by high-performance liquid chromatography 589(1992)346

Arrio, B., see Hill, M. 589(1992)101

Athalye, A. M., Gibbs, S. J. and Lightfoot, E. N.
Predictability of chromatographic protein separations. Study of size-exclusion media with narrow particle size distributions 589(1992)71

Baeseler, M., Boeden, H.-F., Koelsch, R. and Lasch, J.
Cellulose beads: a weak leaking affinity support 589(1992)93

Beard, A., Naikwadi, K. and Karasek, F. W.
Comparison of extraction methods for polychlorinated dibenzo-*p*-dioxins and dibenzofurans in fly ash using gas chromatography–mass spectrometry 589(1992)265

Boeden, H.-F., see Baeseler, M. 589(1992)93

Borrull, F., see Calull, M. 589(1992)151

Brinkman, U. A. T., see Farjam, A. 589(1992)141

Brown, M. A., see Kim, I. S. 589(1991)177

Calull, M., López, E., Marcé, R. M., Olucha, J. C. and Borrull, F.
Optimization of an ion-exchange high-performance liquid chromatographic method for the determination of carboxylic acids, sugars, glycerol and ethanol in wines 589(1992)151

Carr, M. R., see Revill, A. T. 589(1992)281

Caslavska, J., see Thormann, W. 589(1992)321

Chawla, B. and Di Sanzo, F.
Determination of sulfur components in light petroleum streams by high-resolution gas chromatography with chemiluminescence detection 589(1992)271

Chmelík, J., see Thormann, W. 589(1992)321

Collier, P. D., see Van Bergen, C. A. 589(1992)109

Conti, P., see Hamoir, T. 589(1992)31

Craft, N. E., Wise, S. A. and Soares, Jr., J. H.
Optimization of an isocratic high-performance liquid chromatographic separation of carotenoids 589(1992)171

Cromie, D. D. O., see Van Bergen, C. A. 589(1992)109

D'Agostino, P. A. and Provost, L. R.
Determination of chemical warfare agents, their hydrolysis products and related compounds in soil 589(1992)287

Delgado, C., see Tejedo, M. C. 589(1992)127

Desbène, P.-L., see Oliveros, L. 589(1992)53

Desmazières, B., see Oliveros, L. 589(1992)53

De Smet, M., see Hamoir, T. 589(1992)31

Di Sanzo, F., see Chawla, B. 589(1992)271

Dunn, D. L., see Sternitzke, K. D. 589(1992)159

Eisenbeiss, A., Reuke, S. and Türck, M.
Determination of diacetoneketogulonic acid in water samples by high-performance thin-layer chromatography 589(1992)390

Ezra, I., see Massil, S. E. 589(1992)375

Fan, T. Y., see Sternitzke, K. D. 589(1992)159

Fang, X.-H., see Zeng, Z.-R. 589(1992)368

Farjam, A., Van de Merbel, N. C., Nieman, A. A., Lingeman, H. and Brinkman, U. A. T.
Determination of aflatoxin M1 using a dialysis-based immunoaffinity sample pretreatment system coupled on-line to liquid chromatography. Reusable immunoaffinity columns 589(1992)141

Fu, C., see Wang, Z. 589(1992)349

Fuss, L., see Massil, S. E. 589(1992)375

Galantino, M., see Rusconi, L. 589(1992)362

Gibbs, S. J., see Athalye, A. M. 589(1992)71

Głód, B. K., Haddad, P. R. and Alexander, P. W.
Potentiometric detection of carboxylic acids and inorganic anions in ion-exclusion chromatography using camphorsulphonic acid as eluent 589(1992)209

Głód, B. K., see Alexander, P. W. 589(1992)201

Grupeli, M., see Tejedo, M. C. 589(1992)127

Guiochon, G., see Roles, J. 589(1992)223

Haddad, P. R., see Alexander, P. W. 589(1992)201

Haddad, P. R., see Głód, B. K. 589(1992)209

Hague, G. R., see Mulholland, F. 589(1992)380

Haldna, Ü.
Carbonate eluents in ion chromatography. Fractions of resin capacity linked to anions present in the eluent 589(1992)197

Hamoir, T., De Smet, M., Pyrins, H., Conti, P., Vanden Driessche, N., Massart, D. L., Maris, F., Hindriks, H. and Schoenmakers, P. J.
Feasibility study for the construction of an integrated expert system in high-performance liquid chromatography 589(1992)31

Hartwick, R. A., see Wang, T. 589(1992)307

Hill, M. and Arrio, B.
Activation of matrices by 4,6-diphenylthieno[3,4-*d*]-1,3-dioxol-2-one-5,5-dioxide. High-performance liquid affinity chromatographic separations 589(1992)101

Hindriks, H., see Hamoir, T. 589(1992)31

Hirokawa, T., see Hu, J.-Y. 589(1992)339

Horváth, C., see Hung-Jye Lin, F. 589(1992)185

Hu, J.-Y., Hirokawa, T., Nishiyama, F. and Kiso, Y.
Study of isotachophoretic separation behaviour of metal cations by means of particle-induced X-ray emission. II. Separation efficiency of lanthanoid model mixture 589(1992)339

Hua, R. L., see Peng, C. T. 589(1992)231

Huang, Z.-F., see Zeng, Z.-R. 589(1992)368
Hudson, M. K., see Mofidi, M. 589(1992)241
Hung-Jye Lin, F. and Horváth, C.
Measurement of stability constants for complexes of divalent cations with tartaric acid by ion chromatography 589(1992)185
Jeng, C.-Y. and Langer, S. H.
Reaction kinetics and kinetic processes in modern liquid chromatographic reactors (Review) 589(1992)1
Jozefowicz, M., see Letourneur, D. 589(1992)87
Just, U., see Pasch, H. 589(1992)295
Kamiński, M.
Simple test for determination of the degree of distortion of the liquid-phase flow profile in columns for preparative liquid chromatography 589(1992)61
Kane, R. E., see Shansky, R. E. 589(1992)165
Kaniansky, D., see Madajová, V. 589(1992)329
Karasek, F. W., see Beard, A. 589(1992)265
Khoo, S. B., see Yik, Y. F. 589(1992)333
Kim, I. S., Sasinos, F. I., Rishi, D. K., Stephens, R. D. and Brown, M. A.
Determination of aromatic sulfonic acids in aqueous environmental samples by anion-exchange chromatography coupled to particle beam mass spectrometry and UV spectrophotometry 589(1991)177
Kiso, Y., see Hu, J.-Y. 589(1992)339
Koelsch, R., see Baeseler, M. 589(1992)93
Kovács-Hadady, K.
Study of the retention behaviour of barbiturates by overpressured layer chromatography using silica gel bonded with tricaprylmethylammonium chloride 589(1992)301
Krüger, H., see Pasch, H. 589(1992)295
Kunkel, M. E., Seo, A. and Thomas, R. L.
Determination of profiles of non-collagenous proteins from rat bones by sodium dodecyl sulfate high-performance liquid chromatography 589(1992)135
Langer, S. H., see Jeng, C.-Y. 589(1992)1
Lasch, J., see Baeseler, M. 589(1992)93
Lee, H. K., see Yik, Y. F. 589(1992)333
Letourneur, D., Migonney, V., Muller, D. and Jozefowicz, M.
Phosphorylated polystyrene resins in high-performance ion-exchange chromatography 589(1992)87
Li, S. F. Y., see Yik, Y. F. 589(1992)333
Lightfoot, E. N., see Athalye, A. M. 589(1992)71
Lingeman, H., see Farjam, A. 589(1992)141
Liu, D.-M.
Gas chromatographic separation of diastereomeric amino acid derivatives on chiral stationary phases. Application to the determination of enantiomeric composition in (*S*)-(+)-2-butanol 589(1992)249
Liu, W., see Zhou, W. 589(1992)358
López, E., see Calull, M. 589(1992)151
López-Barea, J., see Alhama, J. 589(1992)121
Lucas, R. A., see Van Bergen, C. A. 589(1992)109
Luque, J., see Tejedo, M. C. 589(1992)127
Madajová, V., Turcelová, E. and Kaniansky, D.
Influence of poly(vinylpyrrolidone) on isotachophoretic separations of inorganic anions in aqueous electrolyte systems 589(1992)329
Maltby, D., see Peng, C. T. 589(1992)231
Marcé, R. M., see Calull, M. 589(1992)151
Maris, F., see Hamoir, T. 589(1992)31
Massart, D. L., see Hamoir, T. 589(1992)31
Massil, S. E., Ezra, I. and Fuss, L.
Direct gas chromatographic analysis of O,O-dimethyl- and O,O-diethyldithiophosphoric acids and identification of their by-products 589(1992)375
Migonney, V., see Letourneur, D. 589(1992)87
Minguillón, C., see Oliveros, L. 589(1992)53
Mofidi, M., Hudson, M. K. and Underhill, K.
Flame infrared emission detection of hydrogen 589(1992)241
Molteni, S., see Thormann, W. 589(1992)321
Much, H., see Pasch, H. 589(1992)295
Mulholland, F. and Hague, G. R.
Monitoring 4-methoxy-2,3,6-trimethylbenzenesulphonyl deprotection of arginine-containing synthetic peptides using capillary zone electrophoresis 589(1992)380
Muller, D., see Letourneur, D. 589(1992)87
Naikwadi, K., see Beard, A. 589(1992)265
Neff, W. E., Zeitoun, M. A. M. and Weisleder, D.
Resolution of lipolysis mixtures from soybean oil by a solid-phase extraction procedure 589(1992)353
Nieman, A. A., see Farjam, A. 589(1992)141
Nishiyama, F., see Hu, J.-Y. 589(1992)339
Oliveros, L., Minguillón, C., Desmazières, B. and Desbène, P.-L.
Silica-bonded chiral stationary phases with structurally simple π-donor chiral selectors for high-performance liquid chromatography 589(1992)53
Olucha, J. C., see Calull, M. 589(1992)151
Ong, C. P., see Yik, Y. F. 589(1992)333
Pasch, H., Krüger, H., Much, H. and Just, U.
Analysis of benzyloxy-terminated poly(1,3,6-trioxocane)s by supercritical fluid chromatography 589(1992)295
Peng, C. T., Hua, R. L. and Maltby, D.
Prediction of retention indexes. IV. Chain branching in alkylbenzene isomers with C_{10-13} alkyl chains identified in a scintillator solvent 589(1992)231
Petersson, G., see Strömvall, A.-M. 589(1992)385
Pirkle, W. H. and Welch, C. J.
Effect of superfluous remote polar functionality on chiral recognition 589(1992)45
Preston, H. D., see Van Bergen, C. A. 589(1992)109
Provost, L. R., see D'Agostino, P. A. 589(1992)287
Pyrins, H., see Hamoir, T. 589(1992)31
Reuke, S., see Eisenbeiss, A. 589(1992)390
Revill, A. T., Carr, M. R. and Rowland, S. J.
Use of oxidative degradation followed by capillary gas chromatography–mass spectrometry and multi-dimensional scaling analysis to fingerprint unresolved complex mixtures of hydrocarbons 589(1992)281
Rishi, D. K., see Kim, I. S. 589(1991)177
Roldán, J. M., see Alhama, J. 589(1992)121
Roles, J. and Guiochon, G.
Prediction of the elution profiles of high-concentration bands in gas chromatography 589(1992)223

Roman, M. C. and Siergiej, R. W.
Accuracy of computer-simulated gas chromatographic separations based on a linear elution strength model 589(1992)215
Rowland, S. J., see Revill, A. T. 589(1992)281
Rusconi, L. and Galantino, M.
Solvent system for the counter-current purification of hydrophobic bombesin analogues 589(1992)362
Sasinos, F. I., see Kim, I. S. 589(1991)177
Schoenmakers, P. J., see Hamoir, T. 589(1992)31
Seo, A., see Kunkel, M. E. 589(1992)135
Shansky, R. E. and Kane, R. E.
Separation of soy lecithin using gel permeation chromatography 589(1992)165
Siergiej, R. W., see Roman, M. C. 589(1992)215
Sissons, D. J., see Van Bergen, C. A. 589(1992)109
Soares, Jr., J. H., see Craft, N. E. 589(1992)171
Søeberg, H., see Vinther, A. 589(1992)315
Stephens, R. D., see Kim, I. S. 589(1991)177
Sternitzke, K. D., Fan, T. Y. and Dunn, D. L.
High-performance liquid chromatographic determination of pilocarpine hydrochloride and its degradation products using a β-cyclodextrin column 589(1992)159
Stoev, G.
Application and chiral recognition of heptakis (2,6-di-O-methyl-3-O-trifluoroacetyl)-β-cyclodextrin as a stationary phase for the gas chromatographic separation of enantiomers 589(1992)257
Strömvall, A.-M. and Petersson, G.
Protection of terpenes against oxidative and acid decomposition on adsorbent cartridges 589(1992)385
Tejedo, M. C., Delgado, C., Grupeli, M. and Luque, J.
Affinity partitioning of erythrocytic phosphofructokinase in aqueous two-phase systems containing poly(ethylene glycol)-bound Cibacron Blue. Influence of pH, ionic strength and substrates/effectors 589(1992)127
Thomas, R. L., see Kunkel, M. E. 589(1992)135
Thormann, W., Caslavska, J., Molteni, S. and Chmelík, J.
Capillary isoelectric focusing with electroosmotic zone displacement and on-column multichannel detection 589(1992)321
Tomba, C., see Anelli, P. L. 589(1992)346
Toribio, F., see Alhama, J. 589(1992)121
Turcelová, E., see Madajová, V. 589(1992)329
Türck, M., see Eisenbeiss, A. 589(1992)390
Uggeri, F., see Anelli, P. L. 589(1992)346
Underhill, K., see Mofidi, M. 589(1992)241
Van Bergen, C. A., Collier, P. D., Cromie, D. D. O., Lucas, R. A., Preston, H. D. and Sissons, D. J.
Determination of chloropropanols in protein hydrolysates 589(1992)109
Van de Merbel, N. C., see Farjam, A. 589(1992)141
Vanden Driessche, N., see Hamoir, T. 589(1992)31
Vinther, A. and Søeberg, H.
Radial pH distribution during capillary electrophoresis with electroosmotic flow. Analysis with high ionic strength buffers 589(1992)315
Wang, T. and Hartwick, R. A.
Binary buffers for indirect absorption detection in capillary zone electrophoresis 589(1992)307
Wang, Y.-T., see Zeng, Z.-R. 589(1992)368
Wang, Z., Xu, H. and Fu, C.
Sensitive fluorescence detection of some nitrosamines by precolumn derivatization with dansyl chloride and high-performance liquid chromatography 589(1992)349
Weisleder, D., see Neff, W. E. 589(1992)353
Welch, C. J., see Pirkle, W. H. 589(1992)45
Wise, S. A., see Craft, N. E. 589(1992)171
Wu, C.-Y., see Zeng, Z.-R. 589(1992)368
Xu, H., see Wang, Z. 589(1992)349
Yik, Y. F., Ong, C. P., Khoo, S. B., Lee, H. K. and Li, S. F. Y.
Separation of polycyclic aromatic hydrocarbons by micellar electrokinetic chromatography with cyclodextrins as modifiers 589(1992)333
Zeitoun, M. A. M., see Neff, W. E. 589(1992)353
Zeng, Z.-R., Wu, C.-Y., Fang, X.-H., Huang, Z.-F. and Wang, Y.-T.
Study of dihydroxy-substituted saturated urushiol crown ether as a stationary phase in capillary gas chromatography 589(1992)368
Zhou, W., Liu, W. and An, D.
Simultaneous ion chromatographic determination of chloride and calcium contained in Ringer's injection using a chelating agent as eluent 589(1992)358